

## Colonoscopy Procedure

How to prepare for your Colonoscopy using:

**Patient Name:** 

## **Miralax Split Bowel Prep**

If you have any questions, please call 802-447-5551, Mon.-Fri., 8 a.m.- 3 p.m. If it is after hours, please call the switchboard at 802-442-6361 and request a return call from the Endoscopy nurse on call.

Colonoscopy scheduled with Dr
Date of procedure:Arrival time:
*Follow all of the instructions in this handout exactly as they are written. If you do not follow the directions, your colonoscopy may be cancelled.
*Check with your health insurance provider regarding your health plan benefits prior to your procedure.
*Your bowel must be empty so that your doctor can clearly view your colon.
*It is required for you to have a driver at the time of your discharge.
* Pre-Registration will call you a week prior to your procedure (Arrival times are subject to change.)
*If your procedure is being done under monitored Anesthesia care (heavy sedation), you will also receive a separate call from a preop nurse a week prior to your procedure.
* Failure to complete pre-registration/preop nurse call, will result in your procedure being cancelled.
Instructions on Diabetic and/or Blood Thinners:
Diabetic:
BloodThinners:
***Aspirin, Advil, Aleve, Acetaminophen or Tylenol are acceptable.

Revised 11/16/2023

Continued on next page→

\*\*1 Bottle of Miralax (238 grams)

\*\*1 box of Dulcolax Laxative TABLETS- 5 mg each

## **DAY BEFORE PROCEDURE**

### **NO FOOD, Clear Liquid Diet ONLY**

See clear liquid diet sheet enclosed

#### **BEGIN BOWEL PREP:**

- \*\*Drink 2 quarts of clear liquids throughout the day----Stay hydrated for best results.
- \*\* Anytime Between 1-3 pm, take 2 Dulcolax tablets.
- \*\* <u>Anytime Between 3-5 pm</u>, mix entire 238-gram bottle of Miralax prep into one 64-ounce container
  - or two 32-ounce containers of clear liquid (see enclosed clear liquid sheet for substitutions.)
- \*\* Drink ½ of the mixture (32 oz.). Drink this over a 1 hour period. (8 oz. every 15-30 minutes works well.
- \*\*Continue to drink as much clear liquid as possible until bedtime.
- -If nausea or vomiting develops, stop drinking and wait 30 minutes before restarting.
- -Unless otherwise instructed, please take your regular evening medications after you complete the first half of your prep.

# DAY OF PROCEDURE NOTHING TO DRINK except:

- \*\*Drink the remaining half of the prepared Miralax mixture (32 oz.) by
- \*\*If nausea or vomiting develops, stop drinking and wait 30 minutes before restarting. After Completing your Prep:
  - \*\*Take your normal morning medication with a sip of water.
    - Do not take your diuretic (water pill) this morning.
    - Follow special instructions on **Diabetic/Blood thinner** medications from previous page.
    - -After taking your medications as instructed, nothing further by mouth prior to your appointment.
  - \*\*Take inhalers as prescribed and bring them with you to your appointment.

Please have all jewelry removed for the day of your exam.